

Thresholds found for unilateral optic nerve lesions in MS

17 May 2019



differences in acuities and greater intereye retinal layer thickness differences.

"These thresholds may be useful in establishing the presence of asymptomatic and symptomatic optic nerve lesions in multiple [sclerosis](#) and could be useful in a new version of the diagnostic criteria," the authors write.

More information: [Abstract/Full Text](#)
([subscription or payment may be required](#))

Copyright © 2019 [HealthDay](#). All rights reserved.

(HealthDay)—A new anatomic threshold may be useful for identifying unilateral optic nerve lesions in patients with multiple sclerosis, according to a study published in the May issue of the *Annals of Neurology*.

Rachel C. Nolan-Kenney, from the New York University School of Medicine in New York City, and colleagues measured [optical coherence tomography](#) for 1,530 [patients](#) with multiple sclerosis and healthy controls seen at 11 international sites as part of the International Multiple Sclerosis Visual System Consortium. For this study, presence of an optic nerve lesion was defined as history of acute unilateral optic neuritis.

The researchers found that for identifying unilateral optic neuritis, receiver operating characteristic curve analysis showed an optimal peripapillary retinal nerve fiber layer intereye difference threshold of 5 μm and ganglion cell + inner plexiform layer threshold of 4 μm (477 patients). An association was noted between greater intereye

APA citation: Thresholds found for unilateral optic nerve lesions in MS (2019, May 17) retrieved 14 June 2021 from <https://medicalxpress.com/news/2019-05-thresholds-unilateral-optic-nerve-lesions.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.