


High levels of rare gut bacteria may be linked to restless legs syndrome

7 June 2019



Credit: CC0 Public Domain

Small intestine bacterial overgrowth (SIBO) may be more prevalent among patients with restless legs syndrome (RLS), according to preliminary findings from a small, new study.

Results show that SIBO was found in all seven participants who have RLS. In contrast, the prevalence of SIBO in the [general population](#) is estimated to be no more than 15%.

"We've observed extremely high rates of small intestinal bacterial overgrowth in the RLS group," said lead author Daniel Jin Blum, Ph.D., D.B.S.M., an adjunct clinical instructor at Stanford Center for Sleep Sciences and Medicine in Redwood City, California. "Exploring the relationship between RLS and gut microbial health has the potential to open novel avenues for possible detection, prevention

and treatment for RLS and other [sleep disorders](#)."

SIBO is a condition in which rare gut-residing bacteria are over-represented in the gut. [RLS](#) is a sensorimotor disorder characterized by a complaint of a strong, nearly irresistible urge to move the limbs that is often accompanied by other uncomfortable sensations. These symptoms begin or worsen during periods of rest or inactivity such as lying down or sitting, are partially or totally relieved by movement such as walking or stretching, and occur exclusively or predominantly in the evening or at night.

Low iron in the brain is a key risk factor for RLS. According to the authors, this brain iron deficiency may be secondary to dietary iron deficiency or, potentially, gut inflammation.

Study participants completed questionnaires concerning sleep and SIBO symptoms and took home a fecal collection kit and a SIBO breath test kit. Fecal samples were examined by the University of Minnesota Genomics Center, and SIBO breath samples were evaluated by Aerodiagnostics for hydrogen and methane abnormalities.

Additional study participants continue to be recruited at the Stanford Sleep Center. Further analyses will examine fecal microbial composition, subtypes of RLS [iron](#) deficiency, and comparisons with insomnia.

More information: Daniel J Blum et al, 0009 Restless Leg Syndrome: Does It Start With A Gut Feeling?, *Sleep* (2019). [DOI: 10.1093/sleep/zsz067.008](#)

Provided by American Academy of Sleep Medicine

APA citation: High levels of rare gut bacteria may be linked to restless legs syndrome (2019, June 7)
retrieved 15 June 2021 from <https://medicalxpress.com/news/2019-06-high-rare-gut-bacteria-linked.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.