

Americans still eat too much processed meat and too little fish

June 21 2019



Credit: CC0 Public Domain

A new study in the *Journal of the Academy of Nutrition and Dietetics* found that the amount of processed meat consumed by Americans has remained unchanged in the past 18 years, nor has their intake of fish/shellfish increased. In addition, one quarter of US adults are still eating more unprocessed red meat than the recommended level, and less than 15 percent meet the guidelines for fish/shellfish consumption. On a positive note, Americans are eating less beef and more chicken than they

did 18 years ago, and in fact, for the first time, consumption of poultry exceeds that of unprocessed red meat.

"Despite strong evidence linking processed meat with [cancer risk](#), [consumption](#) of processed meat among US adults didn't change over the study period (1999-2016)," said lead investigator Fang Fang Zhang, MD, Ph.D., Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, USA. "While factors other than health (e.g., social, cultural, and economic) can influence Americans' food choices, the lack of widespread awareness of health risks associated with processed meat may have contributed to the lack of consumption change in the past 18 years. Our findings support further actions to increase the public awareness of the health risks associated with high processed [meat consumption](#) in the US."

The study used a nationally representative sample comprised of dietary data from nearly 44,000 US adults (ages 20 and older) who participated in the National Health and Nutrition Examination Survey (NHANES), through 2016. The investigators assessed trends in consumption of processed meat, [unprocessed red meat](#), poultry, fish, and shellfish over the past 18 years and their purchase locations.

In addition to the overall trends noted above using full NHANES data, the research team also compared NHANES data from 1999-2000 to 2015-2016. Key findings include:

Processed meats: Consumption remained unchanged—182 grams/week compared with 187 grams/week.

- Top five consumed (percentage among total, 2015-2016): Luncheon meat (39 percent), sausage (24 percent), hot dog (9 percent), ham (9 percent), and bacon (5 percent)
- Primary purchase locations: Stores and fast-food restaurants

Unprocessed red meat:

Decreasing trend—340 grams/week compared with 284 grams/week, primarily due to decreased consumption of beef (down by 78 grams/week).

Poultry: Increasing trend—256 grams/week compared with 303 grams/week, primarily due to increased consumption of chicken (up by 34 grams/week)

Fish/seafood: Consumption remained unchanged—115 grams/week compared with 116 grams/week

There is accumulating evidence linking excessive consumption of processed meat to increased risk of obesity, diabetes, cardiovascular diseases, and some cancers. Processed meat has been classified as "carcinogenic to humans" (Group 1) by the International Agency for Research on Cancer (IARC). The American Cancer Society (ACS), the World Cancer Research Fund (WCRF)/American Institute for Cancer Research (AICR) issued recommendations limiting processed meat consumption for cancer prevention. A [study](#) by Dr. Zhang published last month estimated that 14,524 new cancer cases were attributable to high consumption of processed meat in 2015 among US adults aged 20 years and older. Future research is needed to identify barriers to reducing processed meat consumption, evaluate the effectiveness of potential public health interventions, and explore policies such as nutrition quality standards, excise taxes, and health warning labels.

The low consumption of fish/shellfish among US adults could be due to its high retail price, lack of awareness of its health benefits, and concerns about mercury contamination in certain fish, although the scientific evidence suggests that the benefits of fish intake exceed the potential risks for most individuals. Given that fish consumption

(2015-2016) was only half of the recommended level in the 2015-2020 Dietary Guidelines for Americans, efforts are needed to promote the consumption and variety of seafood, especially those varieties high in omega-3 fatty acids.

"Findings of this study can inform public health policy priorities for improving diet and reducing chronic disease burden in the US. Because stores and fast-food restaurants are main purchase locations for processed meat, future policies may prioritize these as primary sites of intervention for reducing processed [meat](#) consumption among US adults," noted Dr. Zhang.

More information: *Journal of the Academy of Nutrition and Dietetics* (2019). [DOI: 10.1016/j.jand.2019.04.004](https://doi.org/10.1016/j.jand.2019.04.004)

Provided by Elsevier

Citation: Americans still eat too much processed meat and too little fish (2019, June 21) retrieved 19 September 2024 from <https://medicalxpress.com/news/2019-06-americans-meat-fish.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--