


# Younger colorectal cancer patients: A missed opportunity for non-emergency diagnoses

7 August 2019



cancer patients under the age of 50 experience delays in primary care leading to emergency diagnoses: a population-based study, *Colorectal Disease* (2019). DOI: [10.1111/codi.14734](https://doi.org/10.1111/codi.14734)

Provided by Wiley

Cancer — Histopathologic image of colonic carcinoid.  
Credit: Wikipedia/CC BY-SA 3.0

In an analysis of information on 10,463 UK patients diagnosed with colorectal cancer from 2006 to 2013, patients under the age of 50 years were more likely to initially experience non-specific symptoms before being referred to cancer specialists.

In the *Colorectal Disease* analysis, young patients were more likely to present with abdominal pain and via an emergency. They also experienced a longer interval between referral to diagnosis by 12.5 days compared with those aged 60 to 69 years.

"Primary care physicians should be made aware of these differences if there is to be a reduction in missed opportunities to prevent emergency diagnoses," the authors wrote.

**More information:** C. S. Arhi et al, *Colorectal*

APA citation: Younger colorectal cancer patients: A missed opportunity for non-emergency diagnoses (2019, August 7) retrieved 16 October 2019 from <https://medicalxpress.com/news/2019-08-younger-colorectal-cancer-patients-opportunity.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*