

Gluten response in coeliac patients could lead to diagnostic test

7 August 2019



Associate professor Jason Tye-Din with patient. Credit: Walter and Eliza Hall Institute of Medical Research

Distinct markers in the blood of people with coeliac disease have been detected within a few hours of gluten being consumed.

The findings address a longstanding mystery about what drives the adverse reaction to gluten in [coeliac disease](#) and could lead to a world-first blood test for diagnosing the [disease](#).


A potential blood-based test would be a vast improvement on the current approach which requires people to consume gluten for a number of weeks, and even months, for the testing to be accurate.

The research, published today in *Science Advances*, involved an international collaboration of the world's leading coeliac disease experts.

The peer-reviewed study included the Walter and Eliza Hall Institute of Medical Research in Australia, University of Oslo in Norway, and in the U.S., Massachusetts General Hospital and University of Chicago. The study was led by

Boston-based biotechnology company ImmusanT Inc.

Associate Professor Jason Tye-Din, head of coeliac research at the Institute and a gastroenterologist at The Royal Melbourne Hospital said work was now underway to explore the development of a simple blood test for coeliac disease.



Associate professor Jason Tye-Din in laboratory. Credit: Walter and Eliza Hall Institute

"For the many people following a [gluten-free diet](#) without a formal diagnosis of coeliac disease, all that might be required is a blood test before, and four hours after, a small meal of gluten.

"This would be a dramatic improvement on the current approach, which requires people to actively consume gluten for at least several weeks before undergoing an invasive procedure to sample the small intestine," Associate Professor Tye-Din said.

Coeliac disease affects approximately 1.4 per cent of people globally, many of whom remain undiagnosed. Symptoms of the disease are caused by a damaging immune response to gluten. After

consuming gluten, patients can experience reactions such as nausea, vomiting, abdominal pain and diarrhoea.

Dr. Bob Anderson, a joint senior author of the paper and the Chief Scientific Officer of ImmusanT Inc., said the new findings could address an important medical need.

"For the first time we have described the inflammatory reaction that patients with coeliac disease experience in the immediate hours after they are exposed to gluten.

"The unpleasant symptoms associated with the disease are linked to an increase in inflammatory molecules in the bloodstream, such as interleukin-2 (IL-2), produced by T cells of the immune system. This response is similar to what happens when an infection is present, however for people with coeliac disease, gluten is the trigger.

"This information underpins a potential new approach to diagnosis that addresses the emerging medical need to identify patients without coeliac disease who may be better served by other treatments for their chronic symptoms," Dr. Anderson said.

Researchers at ImmusanT first discovered the immune markers while assessing [blood samples](#) during the Phase 1 trial of a potential coeliac therapy called Nexvax2. Gastrointestinal symptoms in patients injected with the gluten peptides, particularly nausea and vomiting, correlated with higher levels of IL-2 in their blood.

Subsequent testing showed the consumption of gluten produced the same IL-2 response in people with coeliac disease.

CEO of Coeliac Australia, Ms Michelle Laforest, said the finding would be welcome news for the coeliac community in Australia.

"The potential for a one-off gluten challenge and [blood](#) test could make a significant difference to many thousands of Australians who report sensitivity to [gluten](#) but have been unable to tolerate the current testing approach.

"It is clear this research has the potential to revolutionise the current testing regime for coeliac disease globally," Ms Laforest said.

More information: G. Goel et al., "Cytokine release and gastrointestinal symptoms after gluten challenge in celiac disease," *Science Advances* (2019).

advances.sciencemag.org/content/5/8/eaaw7756

Provided by Walter and Eliza Hall Institute

APA citation: Gluten response in coeliac patients could lead to diagnostic test (2019, August 7) retrieved 26 October 2020 from

<https://medicalxpress.com/news/2019-08-gluten-response-coeliac-patients-diagnostic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.