

# Some pregnant women are exposed to gadolinium in early pregnancy

20 August 2019



Credit: CC0 Public Domain

A small but concerning number of women are exposed to a commonly used MRI contrast agent early in their pregnancy, likely before many of them are aware that they're pregnant, according to a study published in the journal *Radiology*. The results support adherence to effective pregnancy screening measures to help reduce inadvertent exposures to these contrast agents during early pregnancy.

Gadolinium-based contrast agents (GBCAs) are used in as many as 45 percent of MRI exams in the United States to improve the visualization of organs and tissue. Recent evidence shows that trace levels of gadolinium may be retained in the body after the MRI, although the implications of this are not yet understood.

Gadolinium can cross the placenta and enter the fetal circulation. The safety of GBCAs in pregnant [women](#) has not been established, and their use during [pregnancy](#) is not recommended unless essential to the health of the woman or fetus. Available data from cohort studies and case reports have revealed inconsistent findings

regarding the association between gadolinium and adverse fetal outcomes.

To obtain a more precise idea of the prevalence of GBCA exposure among pregnant women, study lead author Steven Bird, Pharm.D., Ph.D., of the U.S. Food and Drug Administration's (FDA) Division of Epidemiology, and colleagues analyzed data on U.S. pregnancies resulting in live births between 2006 and 2017. The data was collected from 16 partners of the FDA's Sentinel System, a program that allows for active surveillance of healthcare data from multiple sources to monitor the safety of regulated [medical products](#).

The data revealed exposures to GBCAs in 5,457 of 4,692,744 live births, a number corresponding to one in 860 pregnancies. Most of the exposures came from contrast MRI examinations of the head, although the number of pelvic and abdominal MRIs were also noteworthy. Almost three-quarters of exposures occurred during the first trimester. The results strongly suggested that inadvertent exposure to GBCAs may occur before pregnancy is recognized.

"Unintended fetal exposures to gadolinium can occur during early pregnancy among women who are not yet aware they are pregnant. Increased attention to existing pregnancy screening measures may help reduce inadvertent exposures to gadolinium contrast," Dr. Bird said.

The FDA has advised all MRI centers to provide a medication guide to outpatients during their first gadolinium [contrast](#) administration, stating that pregnant women and young children may be at increased risk from gadolinium staying in the body.

The researchers suggested several ways radiology imaging centers might avoid inadvertent administration of GBCAs to [pregnant women](#). Among the tools are a safety screening form asking about the potential for pregnancy, direct

questioning of women by radiologic technologists regarding pregnancy, prominently displayed signs asking women to notify radiology staff if they may be pregnant, and pregnancy testing when appropriate.

The FDA is continuing to monitor reports of adverse events associated with gadolinium exposure in utero. While this study was not designed to assess health outcomes of this early exposure for women and/or infants, the FDA is collaborating on and funding a study to evaluate potential risk for stillbirth and other neonatal adverse effects following in utero exposure to [gadolinium](#) in a large group of pregnant Medicaid beneficiaries.

**More information:** "First-Trimester Exposure to Gadolinium-based Contrast Agents: A Utilization Study of 4.6 Million U.S. Pregnancies" *Radiology*, 2019.

Provided by Radiological Society of North America

APA citation: Some pregnant women are exposed to gadolinium in early pregnancy (2019, August 20) retrieved 21 November 2019 from <https://medicalxpress.com/news/2019-08-pregnant-women-exposed-gadolinium-early.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*