


Who you see matters: Stroke patients benefit more from observing their own hand movements during therapy

23 August 2019



(a)



(b)

(a) Top view of the experiment. A tablet monitor was placed over the participant's right forearms on the desk in front of them. (b) Diagrammatic view of the experiment from the left. There is a space to open the hand, which made it easier to imagine the opening-clench hand movement. Credit: Toshihisa Tanaka, TUAT

Japanese scientists at Tokyo University of Agriculture and Technology (TUAT) have found that for stroke patients, observing their own hand movements in a video-assisted therapy—as opposed to someone else's hand—could enhance brain activity and speed up rehabilitation.

Their findings were first made available online in May and published in the July 7, 2019 issue of *IEEE Transactions on Neural Systems and Rehabilitation Engineering*.

A stroke is caused by obstruction of blood flow to the brain and can leave a patient paralyzed. Brain plasticity, where a healthy region of the brain fulfills the function of a damaged region of the brain, is a key factor in the recovery of motor functions caused by stroke. Studies have shown that

sensory stimulation of the neural pathways that control the [sense of touch](#) can promote [brain plasticity](#), essentially rewiring the brain to regain movement and senses.

To promote brain plasticity, [stroke patients](#) may incorporate a technique called [motor imagery](#) in their therapy. Motor imagery allows a participant to mentally simulate a given action by imagining themselves going through the motions of performing that activity. This therapy may be enhanced by a brain-computer interface technology, which detects and records the patients' motor intention while they observe the action of their own [hand](#) or the hand of another person.

"We set out to determine whether it makes a difference if the participant is observing their own hand or that of another person while they're imagining themselves performing the task," said co-author Toshihisa Tanaka, a professor in the Department of Electrical and Electrical Engineering at TUAT in Japan and a researcher at the RIKEN Center for Brain Science and the RIKEN Center for Advanced Intelligent Project.

The researchers monitored [brain activity](#) of 15 healthy right-handed male participants under three different scenarios. In the first scenario, participants were asked to imagine their hand moving in synchrony with hand movements being displayed in a video clip showing their own hand performing the task, together with corresponding voice cues. In the second scenario, they were asked to imagine their hand moving in synchrony with hand movements being displayed on a video clip showing another person's hand performing the task, together with voice cues. In the third scenario, the participants were asked to open and close their hands in response to voice cues only. Using electroencephalography (EEG)—a non-invasive

method of measuring electrical brain activity where electrodes are placed on the scalp —brain activity of the participants was observed as they performed each task.

The team found meaningful differences in EEG measurements when participants were observing their own hand movement and that of another person. The findings suggest that, in order for motor imagery-based therapy to be most effective, video footage of a patient's own hand should be used.

"Visual tasks where a patient observes their own hand movement can be incorporated into brain-computer interface technology used for stroke rehabilitation that estimates a patient's motor intention from variations in [brain](#) activity, as it can give the patient both visual and sense of movement feedback," Tanaka explains.

More information: Hirotaka Nagai et al, Action Observation of Own Hand Movement Enhances Event-Related Desynchronization, *IEEE Transactions on Neural Systems and Rehabilitation Engineering* (2019). [DOI: 10.1109/TNSRE.2019.2919194](#)

Provided by Tokyo University of Agriculture and Technology

APA citation: Who you see matters: Stroke patients benefit more from observing their own hand movements during therapy (2019, August 23) retrieved 17 October 2019 from <https://medicalxpress.com/news/2019-08-patients-benefit-movements-therapy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.