


# Astrocytes and epilepsy

16 September 2019, by Leigh MacMillan



This is an astrocyte, labeled with GFAP (red), Focal Adhesion Kinase (FAK) green, and nuclear stain To-Pro (blue). Credit: Nathan S. Ivey at TNPRC. Via Wikipedia.

The neurodevelopmental disorder Tuberous Sclerosis Complex (TSC) is characterized by often severe epilepsy, along with autism and psychiatric disorders. Astrocytes—star-shaped glial cells that serve multiple functions in the brain—are suspected to play a role in TSC.

Robert Carson, MD, Ph.D., and colleagues proposed that the water channel aquaporin-4 (AQP4), which is important to the functions of astrocytes, contributes to TSC disease pathology.

Reporting in *Neurobiology of Disease*, they found increased expression of AQP4 in cortical brain tissue removed from TSC patients during epilepsy surgery. Using mouse models and cultured astrocytes, they demonstrated that inactivation of the genes *Tsc1* or *Tsc2* (mutations in these genes cause TSC) resulted in increased AQP4

expression.

Increased AQP4 expression has been previously reported in [temporal lobe epilepsy](#), supporting a broader role for this protein in astrocyte dysfunction that leads to seizure activity. AQP4 may represent a novel therapeutic target for the treatment of epilepsy in TSC and perhaps other seizure disorders.

**More information:** Brittany Short et al. Cerebral aquaporin-4 expression is independent of seizures in tuberous sclerosis complex, *Neurobiology of Disease* (2019). DOI: [10.1016/j.nbd.2019.05.003](https://doi.org/10.1016/j.nbd.2019.05.003)

Provided by Vanderbilt University

APA citation: Astrocytes and epilepsy (2019, September 16) retrieved 21 October 2020 from <https://medicalxpress.com/news/2019-09-astrocytes-epilepsy.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*