

Children bullied by friends and siblings are more likely to think about suicide in their early twenties

9 October 2019



Credit: CC0 Public Domain

Depression, self-harm and suicidal ideation are more prominent in adults in their early twenties if they were bullied at home and at school, a study by researchers at the University of Warwick have found. Researchers stress that intervention is needed to educate people in bullying to reduce it.

[Previous studies](#) have identified that sibling bullying has an effect on [mental health](#) in adolescence, however researchers Professor Dieter Wolke and Dr. Slava Dantchev have now found children who were bullied by siblings and friends are more likely to harm themselves.

In the paper, "The Independent and Cumulative Effects of Sibling and Peer Bullying in Childhood on Depression, Anxiety, Suicidal Ideation, and Self-Harm in Adulthood," published in the journal *Frontiers in Psychiatry*, researchers show there is a long shadow thrown by sibling bullying on [self-harm](#), [suicide attempts](#) and depression at 24 years of age.

Using the Children of the 90s study, they were able to show that children who were bullied by siblings had more [mental health issues](#) in adulthood. If they were also bullied by peers this risk increased further.

The participants were asked to self-report bullying when they were 12 years old, whilst depression, anxiety, suicidal ideation and self-harm were assessed at 24 years old.

Of 3,881 youths studied it was found that 31.2% experienced bullying by a sibling. Of those who both became victims and bullied siblings 15.1% were diagnosed with clinical depression, 35.7% experienced [suicidal ideation](#) and 16.1% self-harmed with a further 4.9% with the intent of suicide.

Those who experienced sibling bullying and peer bullying had double the odds of developing [clinical depression](#) and consider suicide.

Dr. Slava Dantchev of the University of Warwick and the University of Vienna said: "This is the first study to show that being bullied by siblings has adverse effects on mental health into adulthood, when the siblings are not living together anymore. Those bullied at home are also more likely to be bullied by peers and have no save space at school or at home. This further increased their torment and affected their mental health"

Professor Dieter Wolke of the Department of Psychology at the University of Warwick comments: "As [sibling bullying](#) often starts when children are young it will be important to educate and help parents to deal and reduce bullying between siblings in early childhood. This is an area which has been completely overlooked in mental health provision and parent support"

More information: Slava Dantchev et al. The Independent and Cumulative Effects of Sibling and Peer Bullying in Childhood on Depression, Anxiety, Suicidal Ideation, and Self-Harm in Adulthood, *Frontiers in Psychiatry* (2019). [DOI: 10.3389/fpsy.2019.00651](https://doi.org/10.3389/fpsy.2019.00651)

Provided by University of Warwick

APA citation: Children bullied by friends and siblings are more likely to think about suicide in their early twenties (2019, October 9) retrieved 19 November 2019 from <https://medicalxpress.com/news/2019-10-children-bullied-friends-siblings-suicide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.