

International study improves traumatic brain injury diagnosis with the use of precision medicine

November 13 2019



Credit: CC0 Public Domain

The research, in which the professor of the Department of Surgery of the University of Valencia Rafael Badenes has taken part, reveals that

the traumatic brain injury (TBI) affects to a greater extent people over 65 years with comorbidities. This contrasts with the fact that, until now, this ailment was related to young men and in good health. The study recommends improving the initial prognosis of this condition, which causes 1.5 million hospital admissions and 57,000 deaths annually in Europe.

The research, published in journal *Neurology*, and in which 32,500 patients from 69 hospitals in 19 countries have participated, identifies accidental falls as a leading cause of [traumatic brain injury](#), compared to the popular belief that links TBI to traffic accidents. The only exception is represented by patients admitted in the [intensive care unit](#) (ICU), who usually have TBI mostly as a result of traffic accidents.

The study differentiates three levels of sick people: those who go to the [emergency room](#), those who are hospitalized and those who enter the ICU. Each of them shows a different degree of severity of traumatic [brain](#) injury and their recoveries are uneven. Experts explain that people in the first two groups are diagnosed with mild TBI while the severity of TBI in the ICU ranges from moderate to severe.

The results of the work reveal that the assignment of severity to each patient determines the treatment and care they receive, but also their subsequent recovery. 83 percent of people admitted in the ICU, 53 percent of those hospitalized and 30 percent of those who came to the emergency room had not achieved a full recovery six months after the accident, which was also reflected in a worse quality of life.

The authors link these deficiencies in [medical care](#) to the traditional TBI classification, which is outdated. Faced with this method, which they consider ineffective, research suggests an approach to precision medicine. "The technique of advanced neuroimaging and blood biomarkers can improve the characterization of the type of damage and

its severity. The differentiation of patients by care procedures offers an innovative perspective," says Professor Badenes.

To the study "Case-mix, care pathways, and outcomes in patients with traumatic brain injury in CENTER-TBI: a European prospective, multicenter, longitudinal, cohort study" European and Israeli medical centers joined. The research follows the treatment they receive and the real-time evolution of more than 4,500 patients affected by TBI. It also analyses the medical history of a second group of about 28,000 sick people, especially the clinical care to which they were subjected and the progress in their state of health. It subsequently compares the results obtained from the two groups.

The conclusions of this research are in line with a previous study, carried out by CENTER-TBI and presented to the European Commission in 2017, which urged bringing precision medicine to the treatment of traumatic brain [injury](#) to lighten the social and human costs of this condition.

More information: Ewout W Steyerberg et al. Case-mix, care pathways, and outcomes in patients with traumatic brain injury in CENTER-TBI: a European prospective, multicentre, longitudinal, cohort study, *The Lancet Neurology* (2019). [DOI: 10.1016/S1474-4422\(19\)30232-7](#)

Provided by Asociacion RUVID

Citation: International study improves traumatic brain injury diagnosis with the use of precision medicine (2019, November 13) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-11-international-traumatic-brain-injury-diagnosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.