


Research shows 'bad cholesterol' is only as unhealthy as its composition

November 18 2019



Space-filling model of the Cholesterol molecule. Credit: RedAndr/Wikipedia

New research at Ohio University shows that a particular subclass of low-density lipoproteins (LDL), also known as "bad cholesterol," is a much better predictor of potential heart attacks than the mere presence of

LDL, which is incorrect more often than not.

The presence of LDL is considered an indicator for the potential risk of [heart](#) attacks or coronary disease, but studies have shown that about 75 percent of patients who suffer heart attacks have [cholesterol levels](#) that don't indicate a high risk for such an event. Research by Ohio University Distinguished Professor Dr. Tadeusz Malinski and researcher Dr. Jiangzhou Hua in Ohio University's Nanomedical Research Laboratory shows that of the three subclasses that comprise LDL, only one causes significant damage.

"Our studies can explain why a correlation of total "bad" cholesterol with a risk of [heart attack](#) is poor and dangerously misleading—it's wrong three quarters of the time," Malinski said. "These national guidelines may seriously underestimate the noxious effects of LDL cholesterol, especially in cases where the content of subclass B in total LDL is high (50% or higher)."

Malinski's team used nanosensors to measure the concentration of nitric oxide and peroxynitrite in endothelium stimulated by LDL subclasses and reported the findings in a study published in the current issue of *International Journal of Nanomedicine*. Subclass B of LDL was found to be the most damaging to endothelial function and can contribute to the development of atherosclerosis. Therefore, it's not the total amount of LDL [cholesterol](#) one has, but rather the concentration of subclass B to the other two, subclass A and subclass I, that should be used to diagnose atherosclerosis and the risk of heart attack.

"Understanding this could lead to improving the accuracy of diagnosis for the evaluation of cardiovascular disease rates," Malinski said.

"Analyzing the mixture of LDL subclasses may provide a parameter-based model for an early medical diagnosis of estimating the risk of cardiovascular disease."

More information: Jiangzhou Hua et al.

Variable Effects Of LDL Subclasses Of Cholesterol On Endothelial Nitric Oxide/Peroxynitrite Balance – The Risks And Clinical Implications For Cardiovascular Disease

, *International Journal of Nanomedicine* (2019). [DOI: 10.2147/IJN.S223524](https://doi.org/10.2147/IJN.S223524)

Provided by Ohio University

Citation: Research shows 'bad cholesterol' is only as unhealthy as its composition (2019, November 18) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-11-bad-cholesterol-unhealthy-composition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.