

# New study confirms American children and teens are consuming significantly less sugary drinks

November 21 2019

---



Credit: CC0 Public Domain

According to a new study in the *American Journal of Preventive Medicine* the share of children and adolescents consuming sugar-

sweetened beverages (SSBs) and the calories they consume from SSBs declined significantly between 2003 and 2014.

This decline in [consumption](#) was found among children and adolescents in all groups studied, including those participating in the Supplemental Nutrition Assistance Program (SNAP), the US federally funded program that provides food assistance to more than 40 million low-income Americans each month—half of whom are children. However, the study demonstrated that even with the decline, current levels remain too high, with 61 percent of all children and 75.6 percent of SNAP recipients still consuming an SSB on a typical day.

"While the observed declines in children's sugar-sweetened [beverage consumption](#) over the past decade are promising, the less favorable trends among children in SNAP suggest the need for more targeted efforts to reduce [sugary drink consumption](#)," explained lead investigator J. Wyatt Koma, Independent Researcher, Washington, DC, USA.

The investigators used nationally representative dietary data for 15,645 children and adolescents (aged 2 to 19 years) from the 2003 to 2014 National Health and Nutrition Examination Surveys (NHANES). They classified children according to self-reported participation in the SNAP program and [household income](#): 27.8 percent were SNAP participants; 15.3 percent were income-eligible but not SNAP participants; 29.7 percent had lower incomes that were ineligible for SNAP; and 27.2 percent had higher incomes that were ineligible for SNAP. The analysis determined the share of children in the various groups who consumed an SSB on a typical day as well as the per capita daily consumption of SSB calories from 2003-2014. SSBs were defined as any non-alcoholic drink with added sugar, including soda, fruit drinks, and flavored milks, consumed on a given day.

This study yielded several significant findings, notably:

- From 2003 to 2014, the share of children consuming an SSB on a typical day declined significantly across all SNAP participation groups, primarily driven by declines in soda consumption. Among children who were SNAP participants, the percentage drinking SSBs declined from 84.2 percent to 75.6 percent and per capita daily consumption of SSB calories declined from 267 to 182 calories.
- In 2014, nearly one in four children who were income-eligible for the SNAP program consumed a fruit drink on any given day (SNAP participants: 24.8 percent; Income-eligible nonparticipants: 23.4 percent).
- The share of SNAP participants consuming a sports/energy drink on any given day tripled from 2003 to 2014 (from 2.6 percent to 8.4 percent).

Although SNAP's success in reducing hunger and food insecurity in the US is well documented, public health attention has more recently turned to the secondary goals of the SNAP program: improving diet quality and preventing obesity among participants.

Senior author Sara N. Bleich, Ph.D., Department of Health Policy and Management, Harvard T.H. Chan School of Public Health, Boston, MA, USA, observed, "Current policy debates are considering whether the diet quality of SNAP participants can be improved by restricting which items can be purchased with SNAP benefits. Our analysis is important for these discussions. While our results confirm that efforts to decrease SSB consumption over the past decade have been successful, they also suggest that the continued surveillance of [children's](#) SSB consumption by beverage type is important, given the consumption trends for sports/energy drinks and non-traditional SSBs like flavored milks. These trends could reduce or eliminate the past decades' achievements limiting SSB consumption."

**More information:** J. Wyatt Koma et al, Sugary Drink Consumption Among Children by Supplemental Nutrition Assistance Program Status, *American Journal of Preventive Medicine* (2019). [DOI: 10.1016/j.amepre.2019.08.033](https://doi.org/10.1016/j.amepre.2019.08.033)

Provided by Elsevier

Citation: New study confirms American children and teens are consuming significantly less sugary drinks (2019, November 21) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-11-american-children-teens-consuming-significantly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.