


# Modulation of proliferation factors in lung adenocarcinoma

11 December 2019



EGFR activated cases compared to pathway markers. Credit: Melanie Haas Kucherlapati - mkucherlapati@partners.org

A recent study published in *Oncotarget* found molecular subtypes based on copy number, DNA methylation, and mRNA expression had variable proliferation levels, the highest correlating with decreased survival.

Significantly, results suggest EGFR over expression and activation are early alterations that likely stall the replication complex through PCNA phosphorylation creating replication stress responsible for DNA damage response and further mutation, but does not promote increased proliferation itself.

Dr. Melanie Haas Kucherlapati from the Department of Genetics at Harvard Medical School in Boston, Massachusetts, USA as well as the Department of Medicine, Division of Genetics at Brigham and Women's Hospital in Boston, Massachusetts, USA said, "It is well established

that cancer is the result of accumulated genetic changes to [tumor suppressor genes](#) or oncogenes, and that these changes lead to uncontrolled cellular proliferation."

This study focuses on genomic and transcriptional changes to proliferation genes across a LUAD cohort created by The Cancer Genome Atlas, previously subtyped by them on the basis of copy number, DNA methylation, and mRNA expression.

The initial part of this study finds that subtype 2, 3, and 6 cases have highest expression of replication components and subtypes 1, 4, and 5 lowest; subtypes with highest expression have decreased survival.

The second part of this study unexpectedly found that levels of EGFR expression overall were inversely proportional to the expression levels of multiple other important proliferation factors across the TCGA LUAD cohort.

Among the group CLPTM1L, PBXIP1, and URGCP like EGFR while showing increased expression over the EGFR cohort, inversely correlated with the expression of multiple key replication proteins over total LUAD. YKT6, KLHL7, FAM220A, and VOPP1 also had increased expression over the EGFR cohort but directly correlated with high expression of multiple [proliferation](#) genes.

The Kucherlapati Research Team concluded that, when EGFR activated cases do metastasize, they target the brain at a much higher rate than non-EGFR activated cases.

**More information:** Melanie Haas Kucherlapati. Modulation of proliferation factors in lung adenocarcinoma with an analysis of the transcriptional consequences of genomic EGFR activation, *Oncotarget* (2019). DOI: [10.18632/oncotarget.27316](https://doi.org/10.18632/oncotarget.27316)

Provided by Impact Journals

APA citation: Modulation of proliferation factors in lung adenocarcinoma (2019, December 11) retrieved 18 September 2020 from <https://medicalxpress.com/news/2019-12-modulation-proliferation-factors-lung-adenocarcinoma.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*