


Researchers identify new therapeutic target for colorectal cancer

27 December 2019



Cancer — Histopathologic image of colonic carcinoid.
Credit: Wikipedia/CC BY-SA 3.0

Researchers at the University of Toronto have identified a key protein that supports the growth of many colorectal cancers. The study, which will be published December 27 in the *Journal of Cell Biology*, reveals that a protein called Importin-11 transports the cancer-causing protein β catenin into the nucleus of colon cancer cells, where it can drive cell proliferation. Inhibiting this transport step could block the growth of most colorectal cancers caused by elevated β catenin levels.

Around 80% of [colorectal cancers](#) are associated with mutations in a gene called *APC* that result in elevated levels of the β catenin protein. This increase in β catenin is followed by the protein's accumulation in the cell nucleus, where it can activate numerous genes that drive [cell proliferation](#) and promote the growth and maintenance of colorectal tumors. But how β catenin enters the [cell nucleus](#) after its levels rise is poorly understood. "Because the [molecular mechanisms](#) underlying β catenin nuclear transport

remain unclear, we set out to identify genes required for continuous β catenin activity in colorectal cancer cells harboring *APC* mutations," says Stephane Angers, a professor in the Department of Pharmaceutical Sciences at the University of Toronto's Leslie Dan Faculty of Pharmacy.

Using CRISPR DNA editing technology, Angers and colleagues, including graduate student Monika Mis, developed a new technique that allowed them to screen the [human genome](#) for genes that support β catenin's activity in colorectal cancer cells after its levels have been elevated by mutations in *APC*. One of the main genes they identified was *IPO11*, which encodes a protein called Importin-11 that is known to be involved in nuclear import.

Angers and colleagues found that Importin-11 binds to β catenin and escorts it into the nucleus of colorectal cancer cells with mutations in *APC*. Removing Importin-11 from these cells prevented β catenin from entering the nucleus and activating its target genes.

The researchers discovered that Importin-11 levels are often elevated in human colorectal cancers. Moreover, removing Importin-11 inhibited the growth of tumors formed by *APC* mutant cancer cells isolated from patients.

"We conclude that Importin-11 is required for the growth of colorectal cancer cells," Angers says. Learning more about how Importin-11 transports β catenin into the nucleus may help researchers develop new therapies that block this process and reduce the growth of colorectal cancers caused by mutations in *APC*.

More information: Mis et al., *Journal of Cell Biology* (2019). DOI: [10.1083/jcb.201903017](https://doi.org/10.1083/jcb.201903017)

Provided by Rockefeller University Press

APA citation: Researchers identify new therapeutic target for colorectal cancer (2019, December 27) retrieved 23 November 2020 from <https://medicalxpress.com/news/2019-12-therapeutic-colorectal-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.