

Vitamin C-B1-steroid combo linked to lower septic shock mortality in kids

January 21 2020

Treating septic shock in children with a combination of intravenous vitamin C, vitamin B1 and hydrocortisone (a commonly used steroid) is associated with lower mortality, according to a study from Ann & Robert H. Lurie Children's Hospital of Chicago. This is the first pediatric study of the safe and relatively inexpensive treatment for septic shock, and the preliminary data supports the promising outcomes seen in adults. Findings were published in the *American Journal of Respiratory and Critical Care Medicine*.

Septic shock is the result of a severe systemic response to infection causing organ failure and dangerously low blood pressure. It is one of the leading causes of death in critically ill children.

"We were surprised and excited to see a substantial reduction in mortality after treating [septic shock](#) in children with a high dose of vitamin C combined with vitamin B1 and hydrocortisone," says lead author Eric Wald, MD, MSCI, critical care physician at Lurie Children's and Associate Professor of Pediatrics at Northwestern University Feinberg School of Medicine. "While based on a [retrospective analysis](#), our results are especially compelling in that they are very similar to the [positive outcomes](#) found in a recent randomized controlled trial of vitamin C treatment for septic shock in adults."

In their retrospective study, Dr. Wald and colleagues matched 43 patients with septic shock who received the vitamin C-B1-hydrocortisone treatment with 43 patients of similar clinical profile

who did not receive it ([control group](#)), and with 43 patients who received hydrocortisone only as adjunctive therapy. They found that while controls had mortality of 28 percent at 30 days, mortality in patients treated with the vitamin C combination protocol dropped to 9 percent in the same period. Treatment with hydrocortisone alone did not improve mortality (30 percent at 30 days). Similar reductions in mortality were seen at 90 days (14 percent with vitamin C protocol vs. 35 percent in controls and 37 percent in the hydrocortisone only group).

"While it is still unclear why vitamin C appears to reduce mortality from septic shock and we need to dig deeper to understand the mechanism, our results are incredibly promising," says Dr. Wald. "We hope to encourage larger, multi-center studies in children with septic shock to confirm our data."

More information: Eric L Wald et al, Hydrocortisone-Ascorbic Acid-Thiamine Use Associated with Lower Mortality in Pediatric Septic Shock, *American Journal of Respiratory and Critical Care Medicine* (2020). [DOI: 10.1164/rccm.201908-1543LE](https://doi.org/10.1164/rccm.201908-1543LE)

Provided by Ann & Robert H. Lurie Children's Hospital of Chicago

Citation: Vitamin C-B1-steroid combo linked to lower septic shock mortality in kids (2020, January 21) retrieved 19 September 2024 from <https://medicalxpress.com/news/2020-01-vitamin-c-b1-steroid-combo-linked-septic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--