


# Time of day affects global brain fluctuations

February 18 2020

---



The negative association between time of day and brain signal fluctuations was strongest in visual and somatosensory regions. Credit: Csaba Orban

As the day progresses, the strength of the brain's global signal fluctuation shows an unexpected decrease, according to a study published on February 18 in the open-access journal *PLOS Biology* by Csaba Orban and a multi-disciplinary team of scientists from the Faculty of

Engineering, Yong Loo Lin School of Medicine and N.1 Institute of Health at the National University of Singapore.

Circadian rhythms govern diverse aspects of physiology including sleep/wake cycles, cognition, gene expression, temperature regulation, and endocrine signaling. But despite the clear influence of [circadian rhythms](#) on physiology, most studies of brain function do not report or consider the impact of time of day on their findings.

To address this gap in knowledge, the team analysed [functional magnetic resonance](#) imaging (fMRI) data of approximately 900 subjects who were scanned between 8 am and 10 pm on two different days as part of the Human Connectome Project (HCP; <http://www.humanconnectomeproject.org/>). Multiple studies have shown that the brain's global signal fluctuates more strongly when one is drowsy (e.g. after insufficient sleep), and fluctuates less when one is more alert (e.g. after coffee). Based on known circadian variation in sleepiness, the authors hypothesized that global signal fluctuation would be lowest in the morning, increase in the mid-afternoon and dip in the early evening.

Instead, they observed a cumulative decrease in global signal fluctuation as the day progressed. This global decrease was most prominent in visual and somatosensory brain regions, which are known for expressing dynamic fluctuations within individuals over time. Across the whole brain, time of day was also associated with marked decreases in resting-state functional connectivity—the correlated activity between different brain regions when no explicit task is being performed.

"We were surprised by the size of the overall time-of-day effects, since the global fMRI signal is affected by many factors and there is substantial variation across individuals. At the [present moment](#) we don't have a good explanation of the directionality of our findings. However, the fact that we also observed slight time-of-day-associated variation in

the breathing patterns of participants suggests that we may also need to consider clues outside of the brain to fully understand these effects," said Csaba Orban, first author of the study.

Based on the findings, the authors recommend that researchers explicitly report the time of day of fMRI scans and other experimental protocols and measurements, as this could help account for between-study variation in results and potentially even failure to replicate findings.

"We hope these findings will motivate fellow neuroscientists to give more consideration to potential effects of time of day on measures of [brain](#) activity, especially in other large-scale studies where subjects are often scanned throughout the day for logistical reasons," said Thomas Yeo, the study's senior author.

**More information:** Csaba Orban et al, Time of day is associated with paradoxical reductions in global signal fluctuation and functional connectivity, *PLOS Biology* (2020). [DOI: 10.1371/journal.pbio.3000602](https://doi.org/10.1371/journal.pbio.3000602)

Provided by Public Library of Science

Citation: Time of day affects global brain fluctuations (2020, February 18) retrieved 24 April 2024 from <https://medicalxpress.com/news/2020-02-day-affects-global-brain-fluctuations.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.