

Why monkeys choose to drink alone

24 February 2020



Credit: CC0 Public Domain

Why do some people almost always drop \$10 in the Salvation Army bucket and others routinely walk by? One answer may be found in an intricate and rhythmic neuronal dance between two specific brain regions, finds a new Yale University study published Feb. 24 in the journal *Nature Neuroscience*.

The biological roots of generosity and selfishness have long fascinated neuroscientists. As [social animals](#), primates depend upon cooperation; yet in times of scarcity or in the quest for status, selfishness often wins out.

Global imaging studies in humans have shown many brain regions seem to be involved in decisions about sharing. Yale's Steve Chang and colleagues decided to focus on [neuronal activity](#) between two specific brain regions of [monkeys](#) faced with a decision about whether or not to share [fruit juice](#) with another monkey.

In one scenario, the monkey could decide to give a drink to a companion or throw it out. In an alternative scenario, they could drink fruit juice

alone or simultaneously share a drink with another monkey.

It turns out monkeys like to drink alone. But, if the alternative is to see the drink dumped in a bin, they prefer to give the other monkey a juice break

In both scenarios, researchers found distinct patterns of interaction in neuronal activity between the amygdala, a relatively primitive area of the brain, and the medial frontal cortex, an area where more deliberate thoughts originate. When monkeys were generous or pro-social, the interactions between these [brain regions](#) were highly synchronized, occurring at the same rate. When they were being anti-social, this synchronicity was markedly suppressed.

The researchers found that they could use the differences in synchronicity of the interactions to predict what decision the monkey had made: They merely had to look at the neuronal data.

"We found a unique signature of neural synchrony that reflects whether a pro-social or an anti-social decision was made," said Chang, senior author of the *Nature Neuroscience* paper and an assistant professor of psychology and neuroscience.

He and his team also found other key differentiating characteristics in the brain during decision making. For instance, when animals were being pro-social, neuronal interactions were transmitted at one frequency, and when anti-social, at another frequency. The frequency was determined by the [brain](#) region in which neurons fired.

"We all know there are [individual differences](#) in levels of generosity," Chang said. "Maybe Scrooge did not have high levels of synchrony after all."

More information: Specialized medial prefrontal–amygdala coordination in other-regarding decision preference, *Nature Neuroscience* (2020). [DOI: 10.1038/s41593-020-0593-y](#) ,

[nature.com/articles/s41593-020-0593-y](https://www.nature.com/articles/s41593-020-0593-y)

Provided by Yale University

APA citation: Why monkeys choose to drink alone (2020, February 24) retrieved 17 January 2021 from <https://medicalxpress.com/news/2020-02-monkeys.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.