

# Neandertals may have had a lower threshold for pain

23 July 2020



Neandertals and modern humans have mixed and exchanged genes several times over the millennia. Researchers have discovered that people who have inherited a gene variant for an ion channel from Neandertals have a lower pain threshold. Science Photo Library/Daynes, Elisabeth

As several Neandertal genomes of high quality are now available researchers can identify genetic changes that were present in many or all Neandertals, investigate their physiological effects and look into their consequences when they occur in people today. Looking into one gene that carries such changes, Hugo Zeberg, Svante Pääbo and colleagues found that some people, especially from central and south America but also in Europe, have inherited a Neandertal variant of a gene that encodes an ion channel that initiates the sensation of pain.

By using data from a huge population study in the UK, the authors show that people in the UK who carry the Neandertal variant of the ion channel experience more pain. "The biggest factor for how much pain people report is their age. But carrying the Neandertal variant of the ion channel makes you experience more pain similar to if you were

eight years older," says lead author Hugo Zeberg, a researcher at the Max Planck Institute for Evolutionary Anthropology and Karolinska Institutet. "The Neandertal variant of the ion channel carries three amino acid differences to the common, 'modern' variant," explains Zeberg. "While single amino acid substitutions do not affect the function of the ion channel, the full Neandertal variant carrying three amino acid substitutions leads to heightened pain sensitivity in present-day people."

On a [molecular level](#), the Neandertal [ion channel](#) is more easily activated which may explain why people who inherited it have a lowered pain threshold. "Whether Neandertals experienced more pain is difficult to say because pain is also modulated both in the [spinal cord](#) and in the brain," says Pääbo. "But this work shows that their threshold for initiating [pain](#) impulses was lower than in most present-day humans."

**More information:** Hugo Zeberg et al, A Neanderthal Sodium Channel Increases Pain Sensitivity in Present-Day Humans, *Current Biology* (2020). [DOI: 10.1016/j.cub.2020.06.045](https://doi.org/10.1016/j.cub.2020.06.045)

Provided by Max Planck Society

APA citation: Neandertals may have had a lower threshold for pain (2020, July 23) retrieved 20 September 2020 from <https://medicalxpress.com/news/2020-07-neandertals-threshold-pain.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*