

Study finds timing of meals affects body rhythms and metabolic health

July 31 2020, by Abbey Hunton



Credit: CC0 Public Domain

Nearly every cell in the body has its own 24-hour clock, and new research from the Perelman School of Medicine at the University of Pennsylvania shows the way those clocks interact with each other plays a critical role in the health of a person's metabolism. It's widely reported

that shift workers suffer from high rates of obesity and diabetes when their internal clocks do not coordinate with each other, as well as due to irregular eating times. However, little is known about the interaction between internal clocks and eating schedules, and specifically, the impact on overall health.

Now, in a new study published today in *Science*, a team of researchers led by Mitchell Lazar, MD, Ph.D., the Willard and Rhoda Ware, Professor in Diabetes and Metabolic Diseases and director of Penn's Institute for Diabetes, Obesity, and Metabolism, sheds new light on the question.

"The internal clocks in the brain synchronize clocks in [peripheral tissues](#), and misalignment of this system is associated with metabolic dysfunction," said Lazar, the senior author of the study. "But how the environment and [genetic factors](#) control the clocks in peripheral tissue and whether communication exists between clocks in different cell types are largely unknown."

Lazar's team, led by postdoctoral fellow Dongyin Guan, Ph.D., established a new mouse model that can specifically disrupt the internal clock in hepatocytes, the major cell type in the liver, which is the body's metabolic hub. As a result of this disruption, researchers observed an accumulation of triglycerides in the blood that increase the risk of heart disease, diabetes, and stroke. These results indicate the importance of the [internal clocks](#) in peripheral tissue of the liver in maintaining metabolic homeostasis.

Surprisingly, the metabolism of other cell types in the liver were also reprogrammed when the internal clock of hepatocytes was disrupted.

"Our discovery of clock communication between different cell types is very exciting as it suggests a previously unappreciated way that the

body's rhythms are coordinated," said Guan.

Although day/night cycles influence behavioral rhythms, such as sleeping, emerging evidence suggests that [food consumption](#) is an important factor in synchronizing peripheral clocks. Recent research showed time-restricted eating can benefit metabolism, and many dieters try intermittent fasting to lose weight. The Lazar team observed that both food timing and the integrity of the internal clock in the liver altered rhythms of metabolism. Specifically, they showed that nearly half of rhythmic genes are regulated by both the internal clock and when the mice ate their food.

Lazar is hopeful that a better understanding of how food affects the body's internal rhythms could lead to an optimized diet schedule, which could be an important preventive approach for shift workers as well as a potential therapeutic strategy for patients with metabolic disorders such as obesity and diabetes.

More information: Dongyin Guan et al. The hepatocyte clock and feeding control chronophysiology of multiple liver cell types, *Science* (2020). [DOI: 10.1126/science.aba8984](https://doi.org/10.1126/science.aba8984)

Provided by Perelman School of Medicine at the University of Pennsylvania

Citation: Study finds timing of meals affects body rhythms and metabolic health (2020, July 31) retrieved 20 September 2024 from <https://medicalxpress.com/news/2020-07-meals-effects-body-rhythms-metabolic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
