


E-cigarettes can be 'gateway' to cigarettes for teens with no prior intention to smoke

November 9 2020



Credit: CC0 Public Domain

Cigarette smoking remains a leading preventable cause of morbidity and mortality in the United States. And while adolescent cigarette smoking has declined over the past several decades, e-cigarette use presents a new

risk for nicotine use disorder. a new study, published Nov. 9 in the journal *Pediatrics*, finds that e-cigarette use is associated with a higher risk of cigarette smoking among adolescents who had no prior intention of taking up conventional smoking. These findings have strong implications for practice and policy, researchers say.

"Research is showing us that adolescent e-cigarette users who progress to cigarette [smoking](#) are not simply those who would have ended up smoking cigarette anyway," says Olusegun Owotomo, M.D., Ph.D., M.P.H., the study's lead author and a pediatric resident at Children's National Hospital. "Our study shows that e-cigarettes can predispose adolescents to cigarette smoking, even when they have no prior intentions to do so."

In one of the first theory-guided nationally representative studies to identify which adolescent e-cigarette users are at most risk of progressing to cigarette smoking, Researchers looked at data of more than 8,000 U.S. adolescents, ages 12-17, who had never smoked. The data was collected by the [Population Assessment of Tobacco and Health \(PATH\) study](#), an NIH and FDA collaborative nationally representative prospective cohort [study](#) of tobacco use, from 2014-2016. Among adolescents who did not intend to smoke cigarettes in the future, those who used e-cigarettes were more than four times more likely to start smoking cigarettes one year later compared to those who did not use e-cigarettes.

E-cigarette use constitutes a relatively new risk factor for nicotine use disorder among U.S. adolescents. A 2019 study from the Centers for Diseases Control and Prevention found that 28% of [high school students](#) and 11% of [middle school students](#) were current e-cigarette users. With the recent emergence of newer and potentially highly addictive e-cigarette products, adolescents who use e-cigarettes are at increased risk of developing nicotine use disorder and progressing to smoke

conventional cigarettes.

"Abstinence from e-cigarettes can protect teens from becoming future smokers and should be framed as a smoking prevention strategy by all concerned stakeholders," says Dr. Owotomo. "Pediatricians are best positioned to educate patients and families on the clinical and psychosocial consequences of [e-cigarette use](#) and should support education campaigns and advocacy efforts geared to discourage adolescent [e-cigarette](#) use."

Provided by Children's National Hospital

Citation: E-cigarettes can be 'gateway' to cigarettes for teens with no prior intention to smoke (2020, November 9) retrieved 19 September 2024 from <https://medicalxpress.com/news/2020-11-e-cigarettes-gateway-cigarettes-teens-prior.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.