

MMR vaccine could protect against COVID-19

November 20 2020



Credit: CC0 Public Domain

The measles-mumps-rubella (MMR) vaccine has been theorized to provide protection against COVID-19. In a new study published in *mBio*, an open-access journal of the American Society for Microbiology, researchers provide further proof of this by showing that mumps IgG titers, or levels of IgG antibody, are inversely correlated with severity in

recovered COVID-19 patients previously vaccinated with the MMR II vaccine produced by Merck. MMR II contains the Edmonston strain of measles, the Jeryl Lynn (B-level) strain of mumps, and the Wistar RA 27/3 strain of rubella.

"We found a statistically significant inverse correlation between mumps titer levels and COVID-19 severity in people under age 42 who have had MMR II vaccinations," said lead study author Jeffrey E. Gold, president of World Organization, in Watkinsville, Georgia. "This adds to other associations demonstrating that the MMR [vaccine](#) may be protective against COVID-19. It also may explain why children have a much lower COVID-19 case rate than adults, as well as a much lower death rate. The majority of children get their first MMR vaccination around 12 to 15 months of age and a second one from 4 to 6 years of age."

In the new study, the researchers divided 80 subjects into 2 groups. The MMR II group consisted of 50 U.S. born subjects who would primarily have MMR antibodies from the MMR II vaccine. A comparison group of 30 subjects had no record of MMR II vaccinations, and would primarily have MMR antibodies from other sources, including prior measles, mumps, and/or rubella illnesses. The researchers found a significant inverse correlation ($r_s = -0.71$, P

Within the MMR II group, mumps titers of 134 to 300 AU/ml ($n=8$) were only found in those who were functionally immune or asymptomatic. All with mild COVID-19 symptoms had mumps titers below 134 AU/ml ($n=17$). All with moderate symptoms had mumps titers below 75 AU/ml ($n=11$). All who had been hospitalized and required oxygen had mumps titers below 32 AU/ml ($n=5$). Titers were measured by Quest Diagnostics using their standard diagnostic protocol.

"This is the first immunological study to evaluate the relationship between the MMR II vaccine and COVID-19. The statistically

significant inverse correlation between mumps titers and COVID-19 indicates that there is a relationship involved that warrants further investigation," said coauthor David J. Hurley, Ph.D., professor and molecular microbiologist at the University of Georgia. "The MMR II vaccine is considered a safe vaccine with very few side effects. If it has the ultimate benefit of preventing infection from COVID-19, preventing the spread of COVID-19, reducing the severity of it, or a combination of any or all of those, it is a very high reward low risk ratio intervention. Maximum seropositivity is achieved through two vaccinations at least 28 days apart. Based upon our study, it would be prudent to vaccinate those over 40 regardless of whether or not they already have high serum MMR titers."

Provided by American Society for Microbiology

Citation: MMR vaccine could protect against COVID-19 (2020, November 20) retrieved 19 September 2024 from <https://medicalxpress.com/news/2020-11-mmr-vaccine-covid-.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--