

# Earlier menopause, fewer pregnancies linked to early onset of progressive MS

23 November 2020, by From Mayo Clinic News Network, Mayo Clinic News Network



Credit: CC0 Public Domain

Women who experience early menopause may be more likely to face an early onset of progressive multiple sclerosis (MS). That is the finding of a Mayo Clinic study recently published in *Brain Communications*.

Researchers also discovered that the fewer pregnancies a woman had, the more likely an earlier onset of progressive MS. These results highlight the key role sex hormones may play for women with MS.

It is already well-known that multiple sclerosis affects men and women differently. Women are two to three times more likely to be diagnosed with MS, an autoimmune disease in which the body's immune system attacks the protective sheath that covers nerve cells in the brain and spinal cord. Researchers previously had discovered that women are more likely to developing the relapsing-remitting phase of multiple sclerosis at an earlier age than men, and women have more frequent relapses than men. Meanwhile, men's symptoms tend to worsen faster than women, causing them to enter the progressive phase of the disease more

quickly.

Delaying the onset of the progressive phase of MS is important in helping prevent or limit severe disability in the future. That is one of the reasons Mayo Clinic researchers wanted to understand the role women's reproductive histories could play in delaying the onset of progressive MS, says Burcu Zeydan, M.D., a Mayo Clinic researcher and the study's lead author.

The study compared the cases of 137 [postmenopausal women](#) with MS seen at Mayo Clinic to 396 postmenopausal women without MS. To identify participants without MS, the researchers relied on the Rochester Epidemiology Project, a unique medical records linkage system, to gather the necessary patient data. They found that women who underwent menopause before 46 were more likely to experience an [early onset](#) of progressive MS. Meanwhile, pregnancies appeared to have a positive effect when it came to delaying progressive MS.

"There seems to be an association between the number of pregnancies and the onset of progressive MS," Dr. Zeydan says. "The higher the number of pregnancies, the later the progressive MS onset."

That is good news for [women](#) with MS who may have avoided [pregnancy](#) due to concerns about negatively affecting the disease's progression. Indeed, it is the opposite.

Regarding their findings on the association between menopause and progression, Dr. Zeydan cautions that more research is needed to truly understand the potential benefits of perimenopausal hormone therapy. But for patients with MS who are already mulling hormone therapy, this study is worth considering.

"Our findings would be another reason to

encourage patients to use hormone therapy," Dr. Zeydan says.

©2020 Mayo Foundation for Medical Education and Research

Distributed by Tribune Content Agency, LLC

APA citation: Earlier menopause, fewer pregnancies linked to early onset of progressive MS (2020, November 23) retrieved 8 March 2021 from <https://medicalxpress.com/news/2020-11-earlier-menopause-pregnancies-linked-early.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*