


New criteria aid diagnosis of traumatic encephalopathy syndrome

14 April 2021



impairment and/or neurobehavioral dysregulation; and a progressive course. It also requires that these features are not fully accounted for by any other neurologic, psychiatric, or medical condition. Individuals meeting these criteria are graded for functional dependence on five levels, from independent to severe dementia. Based on specific RHI exposure thresholds, core clinical features, functional status, and additional supportive features (e.g., delayed onset, motor signs, and psychiatric features), a provisional level of certainty for CTE pathology is determined.

"The criteria are meant for use in research settings to facilitate investigations into the clinical features associated with CTE pathology and to fill other knowledge gaps, including the development of biomarkers for antemortem diagnosis of CTE," the authors write.

With the support of experts from the Uniformed Services University, diagnostic criteria have been developed for traumatic encephalopathy syndrome (TES), the clinical disorder associated with neuropathologically diagnosed chronic traumatic encephalopathy (CTE); the new consensus diagnostic criteria were published online March 15 in *Neurology*.

Several authors disclosed financial ties to the biopharmaceutical industry; one author served as an expert witness in legal cases involving [brain injury](#) and concussion; one author holds a patent relating to Alzheimer disease.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Douglas I. Katz, M.D., from the Boston University School of Medicine, and colleagues developed evidence-informed, expert consensus research [diagnostic criteria](#) for TES. Evidence was reviewed from all published cases of CTE with neuropathological confirmation; predictive validity data on clinical features in relation to CTE pathology were examined from a large clinicopathologic study (298 cases).

Copyright © 2021 [HealthDay](#). All rights reserved.

The authors note that a diagnosis of TES requires considerable exposure to repetitive head impacts (RHI) from [contact sports](#), [military service](#), or other causes; core clinical features of cognitive

APA citation: New criteria aid diagnosis of traumatic encephalopathy syndrome (2021, April 14) retrieved 20 September 2021 from <https://medicalxpress.com/news/2021-04-criteria-aid-diagnosis-traumatic-encephalopathy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.