

Body mass index, age can affect your risk for neck pain

20 April 2021, by Vandana Suresh



Credit: Unsplash/CC0 Public Domain

With roughly 80% of jobs being sedentary, often requiring several hours of sitting stooped in front of a computer screen, neck pain is a growing occupational hazard. Smartphones and other devices have also caused people to bend their necks for prolonged periods. But is bad posture solely to blame?

In a recent study, researchers at Texas A&M University have found that while poor neck and head postures are indeed the primary determinants of [neck pain](#), body mass index, age and the time of the day also influence the neck's ability to perform sustained or repeated movements.

"Neck pain is one of the leading and fastest-growing causes of disability in the world," said Xudong Zhang, professor in the Wm Michael Barnes '64 Department of Industrial and Systems Engineering. "Our study has pointed to a combination of work and personal factors that strongly influence the strength and endurance of the neck over time. More importantly, since these factors have been identified, they can then be modified so that the neck is in better health and

pain is avoided or deterred."

The results of the study are published online in the journal *Human Factors*, a flagship journal in the field of [human factors](#) and ergonomics.

According to the Global Burden of Disease Study by the Institute for Health Metrics and Evaluation, neck pain is ranked as the fourth leading cause of global disability. One of the main reasons for neck pain has been attributed to lifestyle, particularly when people spend long durations of time with their necks bent forward. However, Zhang said a systematic, quantitative study has been lacking on how personal factors, such as sex, weight, age and work-related habits, can affect neck strength and endurance.

For their experiments, Zhang and his team recruited 20 adult men and 20 adult women with no previous neck-related issues to perform controlled head-neck exertions in a laboratory setting. Instead of asking the participants to hold a specific neck posture for a long time, similar to what might happen at a workplace, they performed "sustained-till exhaustion" head-neck exertions.

"In the laboratory, conducting experiments where subjects do long tasks with their neck can take several hours of data collection, which is not very practical for the experimenters and, of course, the participants in our study," said Zhang. "To solve this problem, our experiments were strategically designed to mimic workplace neck strains but in a shorter period of time."

In these exercises, subjects were seated and asked to put on an augmented helmet that allowed them to exert measurable force by the neck. Then, the researchers asked them to either keep their necks straight or maintain their neck tilted in a forward or backward position. In this position, a force was applied to their head and neck on an adjustable frame. This exertion was either to their maximum

capacity of half of it. Before testing, the researchers noted their subjects' age, body mass index and the time of day.

When Zhang and his team analyzed their data, they found that, as expected, work-related factors like head/neck posture play a very important role in determining both neck strength and endurance. But they also observed that while there was no significant difference between male and female subjects' in neck endurance, body mass index was a significant predictor of neck endurance. Also, to their surprise, the time of day affected the neck's ability to sustain an exertion without fatigue.

"It is intuitive to think that over the course of the day, our necks get more tired since we use it more," Zhang said. "But roughly half of our participants were tested in the morning and the remaining in the afternoon. Also, some of the participants had day jobs and some worked the night shift. Despite this, we consistently found the time-of-day effect on neck endurance."

The researchers said their database of neck strength and endurance is also necessary for building advanced musculoskeletal biomechanical models of the neck, which can then be used to, for example, tease apart specific neck muscles that are more vulnerable to injury.

"Looking ahead, we might have the data to begin evaluating if patients recovering from neck injuries are ready to return to work based on whether their neck strength and [endurance](#) are within the norm," Zhang said. "Also, engineers and designers could utilize our data to make wearable devices, like helmets, that are more ergonomic and less stressful on the [neck](#)."

More information: Suman K. Chowdhury et al, Neck Strength and Endurance and Associated Personal and Work-Related Factors, *Human Factors: The Journal of the Human Factors and Ergonomics Society* (2021). [DOI: 10.1177/0018720820983635](https://doi.org/10.1177/0018720820983635)

APA citation: Body mass index, age can affect your risk for neck pain (2021, April 20) retrieved 12 June 2021 from <https://medicalxpress.com/news/2021-04-body-mass-index-age-affect.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.