


Delaying second dose of COVID-19 vaccines may be an effective public health strategy

April 21 2021



During an outbreak, which vaccination strategy would avert more infections, hospitalizations, deaths: a strategy that provides lower protection to more people, or one that provides higher protection to fewer individuals? Credit: Moghadas (lead author)

Two of the COVID-19 vaccines currently approved in the United States require two doses, administered three to four weeks apart, however, there are few data indicating how best to minimize new infections and

hospitalizations with limited vaccine supply and distribution capacity. A study published on 21st April, 2021 in the open access journal *PLOS Biology* by Seyed Moghadas at York University in Toronto, Canada, and colleagues suggests that delaying the second dose could improve the effectiveness of vaccine programs.

The emergence of novel, more contagious SARS-CoV-2 variants has led to a public health debate on whether to vaccinate more individuals with the first dose of available vaccines and delay the second dose, or to prioritize completion of the two-dose series based on tested schedules in [clinical trials](#).

In order to compare the epidemiological impact of each vaccination strategy, researchers built a [mathematical model](#) that simulated both COVID-19 transmission and various delayed second dose vaccination schedules. The model simulated several scenarios, including a range of levels of preexisting immunity in the population and decreased [vaccine](#) efficacy of the first dose when followed by a longer interval between doses.

The authors found that delaying the second dose for 9-15 weeks after the first dose avoided more hospitalizations, infections, and deaths compared to following the recommended schedules for Moderna and Pfizer-BioNTech vaccines.

The authors note that the study has several limitations, including a lack of clinical evidence quantifying the durability of the vaccines when administered under different schedules. Researchers thus assumed that protection levels of the first dose were stable if second doses were delayed, and that the protection level after delaying second doses was identical to overall protection after two doses when vaccinating according to [schedule](#). Further studies are needed to pinpoint the optimal time between doses for each type of vaccine.

According to the authors, "When racing against a burgeoning outbreak, our results show that prioritizing vaccine coverage with rapid distribution of the first dose would be critical to mitigating adverse outcomes and allow the healthcare system to also address non-COVID-19 medical needs of the population."

"We still do not have the full picture of vaccine effectiveness as new and more contagious variants spread. Efficacy of vaccines against these variants is an additional factor that would need to be considered in determining the outcomes of on-time versus delayed second dose and interval between doses," said Moghadas, the lead author of the study.

More information: Moghadas SM, Vilches TN, Zhang K, Nourbakhsh S, Sah P, Fitzpatrick MC, et al. (2021) Evaluation of COVID-19 vaccination strategies with a delayed second dose. *PLoS Biol* 19(4): e3001211. doi.org/10.1371/journal.pbio.3001211

Provided by Public Library of Science

Citation: Delaying second dose of COVID-19 vaccines may be an effective public health strategy (2021, April 21) retrieved 20 September 2024 from <https://medicalxpress.com/news/2021-04-dose-covid-vaccines-effective-health.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--