


Babies surviving Group B strep more likely to require special educational support

22 April 2021



Streptococcus mutans. Gram stain. Credit: CDC/Public Domain

Invasive Group B Streptococcus (GBS) disease, notably meningitis, during the first days and months of a baby's life can have persistent effects for children and hence their families, according to new research. Published in the *Lancet Child & Adolescent Health*, the study is the first evidence of long-term effects including after GBS sepsis (infection in the bloodstream).

This large study analyzed outcomes for nearly 25,000 children born in Denmark and The Netherlands, between 1997 and 2017. Results show that children who had invasive GBS infection are twice as likely to have neurodevelopmental impairments (NDI) and to require special educational support than children who did not have GBS disease.

Effects later in life are significant—about one in 20 survivors will experience some form of neurodevelopmental disability. The authors note that the majority of GBS infection survivors will not have long-term effects.

This study was conducted through a partnership of the London School of Hygiene & Tropical Medicine (LSHTM), Aarhus University (Denmark), Amsterdam UMC (Netherlands) and the Dutch National Institute for Public Health and the Environment.

Invasive GBS infection is increasingly recognized as a leading cause of newborn illness worldwide, with ~300,000 cases estimated annually. Important knowledge gaps have been highlighted especially regarding the long-term outcomes of patients who survive the acute disease during this critical period of life. Previous small studies have followed babies after GBS meningitis to a median age of about two years. No study has examined outcomes after GBS sepsis.

Dr. Bronner Gonçalves, from LSHTM and a lead author on the study, said: "This is the largest study quantifying long-term effects of GBS infection for babies who survive, with follow-up to the second decade. We add to findings on long-term effects after meningitis, showing subtler cognitive outcomes at older ages. These new findings show an important, previously unquantified risk after GBS sepsis. Our results suggest that continued medical follow-up is key and could lead to earlier identification of problems, and more support to families."

Children in the study were identified using national administrative, medical, and laboratory databases. 2,258 babies were diagnosed as having had GBS infection, including meningitis and sepsis, within the first three months of life. For every GBS case, ten children who did not have the infection were matched on age, sex and other characteristics influencing long-term outcomes.

Children were followed up for an average of fourteen years in Denmark and nine years in the Netherlands, to analyze risks of death and of NDI (e.g. motor, cognitive, and socio-behavioral, and

[special educational needs](#)). The findings are striking including with intrapartum antibiotics becomes more and reinforce the severity of this condition with a 5% mortality risk for babies dying from invasive GBS disease in high-income settings, although the risk is much higher in low and middle income countries. Compellingly, there are also serious consequences for some of the children who survive.

In Denmark, history of GBS infection in the first three months of life was associated with 1.8-fold increase in risk of NDI: 45 of the 969 children who survived the acute phase of the disease had moderate or severe NDI, such as motor impairment, by the age of 10. The risk was even higher in the Netherlands, where 36 of the 252 children with a history of invasive GBS disease needed special educational support by age 10, a more than two-fold risk.

Dr. Erzsébet Horváth-Puhó, an associate professor at Aarhus University, Denmark and a lead author, said: "Our study suggests that babies who recover from invasive GBS disease have increased risks of cognitive, motor, and behavioral impairments, which can impact their lives and their families. The long term effects on children was also evidenced by higher rates of hospitalization for those who survived this infection. In Denmark, a nearly two-fold increase in outpatient visits was observed and a 1.3 higher increase for hospital admissions compared to children who did not develop GBS disease."

Currently, the only available GBS disease prevention strategy is to provide antibiotic prophylaxis to women in labor whose baby is at increased risk of developing GBS disease. However, this strategy is not sufficient even in these two European countries and currently not adopted in settings with limited resources.

Professor Merijn Bijlsma, a pediatrician at Amsterdam University and a lead author, said: "Even with the best medical care, such as available for Dutch and Danish babies, survivors of GBS meningitis and also sepsis may be affected for life. The best way to protect these children is prevention of invasive GBS disease. We need better ways to identify at birth who is at risk, so prevention

Professor Joy Lawn, from LSHTM and senior author and PI, said: "Every year hundreds of thousands of babies have GBS sepsis or meningitis, with a high mortality risk. Families in all continents are affected. Our study shows that GBS also affects educational potential for survivors. The global burden of disease is more than has previously been estimated, underlining the urgent need for prevention. Despite 30 years of research, maternal GBS vaccines currently are not yet available—a contrast to more than 100 COVID-19 vaccines in process within one year. We can and must deliver more for families affected by GBS all over the world."

Charlotte Cleary's daughter Aimee developed early-onset group B Strep meningitis and sepsis shortly after birth in July 2011. She spent three weeks very sick in hospital. and has recovered with brain damage and physical disabilities. Aimee was diagnosed with quadriplegic spastic cerebral palsy and pseudobulbar palsy, along with learning difficulties and sensory issues. Put simply, she can't sit, stand, walk, talk and is mostly tube-fed. Charlotte said: "Every day is a challenge and it's hard to look after a disabled child, although she does bring me so much joy. I do however have anger inside of me that GBS is so easy to prevent, so why are they not doing more to routinely test it in pregnancy? That is my question as Aimee's disability would most likely be non-existent."

The authors acknowledge study limitations including differences in the identification of children with GBS infection cases selection between the two countries. Importantly, these results from Denmark and the Netherlands reflect two high income European countries, and results in low and [middle income countries](#) may differ, hence data are needed from these contexts. Further research is needed as to why different [children](#) are affected differently. We know preterm birth is an important linked risk factor and more studies are needed to explore this association.

More information: Erzsébet Horváth-Puhó et al, Mortality, neurodevelopmental impairments, and

economic outcomes after invasive group B streptococcal disease in early infancy in Denmark and the Netherlands: a national matched cohort study, *The Lancet Child & Adolescent Health* (2021). DOI: [10.1016/S2352-4642\(21\)00022-5](https://doi.org/10.1016/S2352-4642(21)00022-5)

Provided by London School of Hygiene & Tropical Medicine

APA citation: Babies surviving Group B strep more likely to require special educational support (2021, April 22) retrieved 22 June 2021 from <https://medicalxpress.com/news/2021-04-babies-surviving-group-strep-require.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.