

Simple foot test detects heart rhythm disorder in patients with diabetes

24 April 2021



Credit: Unsplash/CC0 Public Domain

Atrial fibrillation can be detected during annual foot assessments in patients with diabetes, according to research presented today at EHRA 2021, an online scientific congress of the European Society of Cardiology (ESC).

"In our study, one in six patients with [diabetes](#) had previously undiagnosed [atrial fibrillation](#)," said study author Dr. Ilias Kanellos of the European University of Cyprus, Nicosia. "This presents an opportunity to provide treatment to prevent subsequent strokes."

Diabetes is an [independent risk factor](#) for atrial fibrillation. Prevalence of the heart rhythm disorder is at least two-fold higher in patients with diabetes compared to those without. Although people with atrial fibrillation have a five times greater risk of stroke, anticoagulation medication is an effective preventive therapy.

Annual foot checks are advised for patients with diabetes. This is because increased blood sugars cause poor circulation, [nerve damage](#) and numbness, and could ultimately lead to foot

amputation if problems go undetected.

Dr. Kanellos said: "The yearly check-up includes palpating the arteries in the feet to examine whether there is healthy blood flow. During this examination we noticed that some patients had an [irregular rhythm](#) and decided to investigate its usefulness for diagnosing atrial fibrillation."

Two podiatrists and six podologists were trained to spot rhythm abnormalities during pulse palpation of the foot arteries. They were also shown how to confirm their findings using a hand-held Doppler ultrasound device.

The 12-month observational study included 300 patients with diabetes attending their annual foot screening appointments. Patients with rhythm abnormalities during pulse palpation of their foot arteries underwent a Doppler ultrasound to verify the observations.

The average age of participants was 60 years. There were 180 men and 120 women. The researchers found that 51 patients (17%) had previously undiagnosed atrial fibrillation. The heart rhythm disorder was found in a similar proportion of men and women. Patients were advised to visit a cardiologist for an electrocardiogram (ECG) to reinforce the diagnosis and discuss treatment.

Dr. Kanellos said: "The study has identified a simple, quick, and low-cost way to identify [patients](#) with diabetes who have a heart rhythm disorder they were unaware of. This intervention has the potential to avoid strokes and preserve quality of life in this patient group."

More information: Abstract title: Routine podiatry assessment as a potential preventive tool for atrial fibrillation screening in diabetics.

Provided by European Society of Cardiology

APA citation: Simple foot test detects heart rhythm disorder in patients with diabetes (2021, April 24) retrieved 21 June 2021 from <https://medicalxpress.com/news/2021-04-simple-foot-heart-rhythm-disorder.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.