

Exoskeleton-assisted walking may improve bowel function in people with spinal cord injury

28 May 2021



Two types of exoskeletons were used in this multi-site study, ReWalk and Ekso GT. This photo shows an Ekso GT in the Tim & Caroline Reynolds Center for Spinal Stimulation at Kessler Foundation. Credit: Kessler Foundation

A team of researchers has shown that physical intervention plans that included exoskeleton-assisted walking helped people with spinal cord injury evacuate more efficiently and improved the consistency of their stool. This finding was reported in *Journal of Clinical Medicine* on March 2, 2021, in the article "The Effect of Exoskeletal-Assisted Walking on Spinal Cord Injury Bowel Function: Results from a Randomized Trial and Comparison to Other Physical Interventions."

The authors are Peter H. Gorman, MD, of the University of Maryland School of Medicine, Gail F. Forrest, Ph.D., of Kessler Foundation's Tim and Caroline Reynolds Center for Spinal Stimulation, Dr. William Scott, of VA Maryland Healthcare System, Pierre K. Asselin, MS, Stephen Kornfeld, MD, Eunyoung Hong, Ph.D., and Ann M. Spungen, EdD, of the James J. Peters VA Medical

Center.

Bowel dysfunction, a [common experience](#) after spinal cord [injury](#), can lead to chronic constipation and incontinence, causing discomfort and frustration. In one survey, more than a third of men with spinal cord injury reported that bowel and bladder dysfunction had the most significant effect on their lives post-injury. Unfortunately, these issues are not easily managed.

Rehabilitation professionals have traditionally managed bowel dysfunction using approaches that target the gastrointestinal system or require manual intervention, but some newer research suggests that physical activity and upright posture may enhance bowel motility. However, few studies have explored the possibility that exoskeletal-assisted walking—in which a person with spinal cord injury wears a robotic suit, enabling them to stand and walk—may be an effective addition to existing intervention plans.

In this study, the research team investigated whether exoskeletal-assisted walking improved bowel function in people with chronic spinal cord injury. They performed a three-center, randomized, controlled, crossover clinical trial in which 50 participants completed 36 sessions of exoskeletal-assisted walking. The researchers evaluated bowel function as a secondary outcome in 49 participants. Bowel function was measured via a 10-question bowel function survey, the Bristol Stool Form Scale, and the Spinal Cord Injury Quality of Life Bowel Management Difficulties instrument.

Results showed that the exoskeletal-assisted walking program provided some improvement in bowel function when compared to a control group. "We saw a notable reduction in bowel evacuation time, with 24 percent of participants reporting an

improved experience," said Dr. Forrest, co-author and associate director of the Center for Mobility and Rehabilitation Engineering Research at Kessler Foundation. "We also noted that participants' stools trended toward better consistency, supporting our hypothesis that this intervention may improve several measures of [bowel](#) function."

"Our results support the idea that walking, and not just standing, may have a beneficial effect on [bowel function](#)," said Dr. Gorman, co-author and chief of the Division of Rehabilitation Medicine at the University of Maryland Rehabilitation and Orthopaedic Institute. "Our goal is to improve the quality of life of those with chronic spinal cord injury, and these encouraging results will help inform future studies on the emerging field of mobility intervention."

More information: Peter H. Gorman et al, The Effect of Exoskeletal-Assisted Walking on Spinal Cord Injury Bowel Function: Results from a Randomized Trial and Comparison to Other Physical Interventions, *Journal of Clinical Medicine* (2021). [DOI: 10.3390/jcm10050964](https://doi.org/10.3390/jcm10050964)

Provided by Kessler Foundation

APA citation: Exoskeleton-assisted walking may improve bowel function in people with spinal cord injury (2021, May 28) retrieved 28 January 2022 from <https://medicalxpress.com/news/2021-05-exoskeleton-assisted-bowel-function-people-spinal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.