


Self-reported declines in cognition may be linked to changes in brain connectivity

22 June 2021



Credit: Pixabay/CC0 Public Domain

Jessica Damoiseaux, Ph.D., an associate professor with the Institute of Gerontology at Wayne State University, recently published the results of a three-year study of cognitive changes in older adults. The team followed 69 primarily African American females, ages 50 to 85, who complained that their cognitive ability was worsening though clinical assessments showed no impairments. Three magnetic resonance imaging scans (MRIs) at 18-month intervals showed significant changes in functional connectivity in two areas of the brain.

"An older adult's perceived [cognitive decline](#) could be an important precursor to dementia," Damoiseaux said. "Brain alterations that underlie the experience of decline could reflect the progression of incipient dementia and may emerge before cognitive assessment is sensitive enough to detect a deficit."

The resulting paper, "Longitudinal change in hippocampal and dorsal anterior insulae [functional connectivity](#) in subjective cognitive decline," appeared in the May 31 issue of *Alzheimer's Research & Therapy*. Damoiseaux conducted the study with graduate student Raymond Viviano,

Ph.D., who is first author.

Subjective cognitive decline, defined as a perceived worsening of cognitive ability not noted on clinical assessment, may be an early indicator of dementia. Previous cross-sectional research has demonstrated aberrant brain functional connectivity in subjective cognitive decline, but longitudinal evaluation has been limited.

Viviano and Damoiseaux's three-year study found that persons reporting more [subjective cognitive decline](#) showed a larger decrease in connectivity between components of the default mode network and a larger increase in connectivity between salience and default mode network components. The functional connectivity changed in the absence of a change in cognitive performance.

Since these brain changes occurred without concomitant cognitive changes, they could indicate that brain changes underlie the perception of decline. These changes could be a sensitive marker for nascent dementia months or years before assessments detect any cognitive deficit.

More information: Raymond P. Viviano et al, Longitudinal change in hippocampal and dorsal anterior insulae functional connectivity in subjective cognitive decline, *Alzheimer's Research & Therapy* (2021). [DOI: 10.1186/s13195-021-00847-y](https://doi.org/10.1186/s13195-021-00847-y)

Provided by Wayne State University

APA citation: Self-reported declines in cognition may be linked to changes in brain connectivity (2021, June 22) retrieved 16 September 2021 from <https://medicalxpress.com/news/2021-06-self-reported-declines-cognition-linked-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.