

Easy detection of fluorescence emitted by protein behind aging

5 August 2021




Figure 1: Fluorescence intensity of young and old *C. elegans*. (Red indicates higher fluorescence intensity.)
Credit: Osaka City University

Professor Emeritus Yoshikazu Nishikawa of Osaka City University's Graduate School of Human Life Science, and collaborators, have found that a substance that emits a specific fluorescence in vivo increases with age. The fluorescence is derived from advanced glycation end products (AGEs) and correlates with the life expectancy of *C. elegans*. The measurement technology of this model could be used for the search and evaluation of anti-glycation materials in the future.

The results of this research were published in the international academic journal *npj Aging and Mechanisms of Disease* on Monday, June 7, 2021.

As the global population ages and [preventive measures](#) against age-related diseases such as dementia and cancer become increasingly important, glycation in the [body](#) is attracting attention as an important risk factor for aging and lifestyle-related diseases. It is known that the amount of AGEs in the body increases with aging and age-related diseases such as diabetes and arteriosclerosis, and measuring the amount of AGEs in the body is expected to be an indicator for

predicting aging and age-related diseases. However, experiments on natural aging using humans and mammals require several years, and AGEs detection methods are mainly invasive blood tests, which require expensive and special experimental operations using antibodies, such as ELISA. The relationship between glycation and aging in the body is still unclear and a simple method for evaluation has not yet been established.

This research group hypothesized that AGEs could be indirectly assessed by measuring the [fluorescence](#) of some AGEs, and succeeded in clarifying the relationship between AGEs and fluorescence in vivo using the model organism *Caenorhabditis elegans*, and in developing a high-throughput evaluation method.




Figure 2: Fluorescence microscope images of young and old *C. elegans* (fluorescent area: blue). Credit: Osaka City University

In this study, the research team first conducted fluorescence spectrum analysis using protein extracts from young and old *C. elegans* and found that specific fluorescence was enhanced in protein extracts from old worms (Fig. 1). They developed a method to detect the fluorescence in vivo and following the same worm over time, they found that

the fluorescence value increased with age. In addition, they found that fluorescence decreased in long-lived *C. elegans* mutants and in ones treated with rifampicin, which has an anti-glycation effect, the lifespan was extended, while fluorescence increased in *C. elegans* treated with ribose, which has a strong glycation effect. Furthermore, when protein extracts from young *C. elegans*, which did not fluoresce, were artificially glycosylated, they fluoresced at the same wavelength as old *C. elegans*, suggesting a strong relationship between this fluorescence, AGEs, and aging. As a result of comparison by analysis of all proteins expressed in old and young worms and observing this with fluorescence microscopy, the team found that this fluorescence was derived from AGEs-derived vitellogenin (Fig. 2).

This suggests that the fluorescence at specific wavelengths emitted from the body of the worm reflects AGEs, and by measuring this fluorescence, the amount of AGEs in the body can be estimated, suggesting the possibility of using it as an indicator of aging.

Since the body of *C. elegans* is transparent, the fluorescence intensity inside the body can be observed while the animal is still alive and they have a life span of about three weeks. In the future, the method laid out in this research is expected to be utilized on *C. elegans* to search for food materials related to anti-glycation and anti-aging.

More information: Tomomi Komura et al, Autofluorescence as a noninvasive biomarker of senescence and advanced glycation end products in *Caenorhabditis elegans*, *npj Aging and Mechanisms of Disease* (2021). DOI: [10.1038/s41514-021-00061-y](https://doi.org/10.1038/s41514-021-00061-y)

Provided by Osaka City University

APA citation: Easy detection of fluorescence emitted by protein behind aging (2021, August 5) retrieved 23 October 2021 from

<https://medicalxpress.com/news/2021-08-easy-fluorescence-emitted-protein-aging.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.