

Smoking exposure during childhood may increase risk of rheumatoid arthritis

August 18 2021



Credit: CC0 Public Domain

A new study by investigators from Brigham and Women's Hospital found a potential direct link between exposure to parental smoking during childhood and increased risk of seropositive rheumatoid arthritis

(RA) later in life. Researchers utilized established longitudinal data from 90,923 women in the Nurses' Health Study II (NHSII) to elucidate the relationship between passive smoking exposure and incident RA. Passive exposure was broken down into three categories, including maternal smoking during pregnancy, parental smoking during childhood, and years lived with smokers since age 18. Even with personal smoking accounted for, passive exposure to parental smoking during childhood was found to increase risk of incident seropositive RA by 75-percent. Findings are published in *Arthritis & Rheumatology*.

"There has been intense interest in mucosal lung inflammation from personal smoking as a site of RA pathogenesis," said senior author Jeffrey A. Sparks, MD, MMSc, of the Department of Medicine at the Brigham. "But the majority of RA patients aren't smokers, so we wanted to look at another inhalant that might precede RA."

RA is an inflammatory disease characterized by arthritis at multiple joints and is associated with morbidity and mortality outcomes. Many people with RA have signs of lung inflammation, and while genetic and [environmental factors](#) contribute to risk of developing RA, smoking has long been implicated as a key RA risk factor. Personal (active) smoking is the most well-established environmental risk factor associated with RA, with passive smoking left relatively unexplored.

To link passive smoking and incident RA more conclusively, Sparks and colleagues used data from NHSII questionnaires collected biennially between 1989 and 2017 from 90,923 women aged 35-52 years. Researchers used participant medical records to confirm incident RA and serostatus. Statistical modeling was then used to estimate the direct effect of each passive smoking exposure on RA risk, as well as to control for other factors such as personal smoking.

A 75-percent higher risk of RA was found in individuals who

experienced passive childhood exposure to parental smoking. This risk increased in participants who themselves became active smokers. Over the median follow-up of 27.7 years, 532 women in the cohort developed confirmed incident RA cases—the majority (352) of which were seropositive (positive for RA autoantibodies). Maternal smoking during pregnancy and years lived with smokers beyond age 18 showed no significant association with incident RA risk.

Although the all-female nurse participant pool led to high response rates and retention, the study is limited by the absence of men. The team intends to continue with longitudinal studies that encompass both men and women, as to provide critical insight into other rheumatoid conditions and even other [autoimmune diseases](#).

"Our findings give more depth and gravity to the negative health consequences of smoking in relation to RA, one of the most common autoimmune diseases," said lead and co-corresponding author Kazuki Yoshida, MD, ScD, of the Brigham's Division of Rheumatology, Inflammation and Immunity. "This relationship between childhood parental [smoking](#) and adult-onset RA may go beyond rheumatology—future studies should investigate whether childhood exposure to inhalants may predispose individuals to general autoimmunity later in life."

More information: Yoshida K et al. "Passive Smoking Throughout the Life Course and the Risk of Incident Rheumatoid Arthritis in Adulthood Among Women" *Arthritis Rheumatol.* [DOI: 10.1002/art.41939](https://doi.org/10.1002/art.41939)

Provided by Brigham and Women's Hospital

Citation: Smoking exposure during childhood may increase risk of rheumatoid arthritis (2021,

August 18) retrieved 20 September 2024 from
<https://medicalxpress.com/news/2021-08-exposure-childhood-rheumatoid-arthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.