


Therapeutic potential of Mozart for medication-resistant epilepsy

16 September 2021



Credit: Pixabay/CC0 Public Domain

Listening to Mozart's Sonata for Two Pianos in D Major (K448) for at least 30 seconds may be associated with less frequent spikes of epilepsy-associated electrical activity in the brain in people with medication-resistant epilepsy. The findings, which also suggest that positive emotional responses to K448 may contribute to its therapeutic effects, are published in *Scientific Reports*.

Previous research has shown that listening to K448 is associated with less frequent spikes of epilepsy-associated electrical activity in the [brain](#) in people with epilepsy. However, the impact of [music](#) duration on this association and the reasons for it have been unclear.

Robert Quon and colleagues used electroencephalogram (EEG) to measure electrical activity in the brains of 16 adults with medication-resistant epilepsy as they listened to a series of either 15 or 90 second music clips, including K448. The authors found that listening to between 30 and 90 seconds of K448, but not the other music clips, was associated with a 66.5% average reduction in

the number of epilepsy-associated electrical activity spikes throughout the brain. These reductions were found to be greatest in the brain's left and right frontal cortices, parts of the brain involved in regulating [emotional responses](#).

The researchers also observed that when participants listened to the ending of long, repetitive sections within K448, a type of electrical activity known as theta activity increased in their frontal cortices. Previous research has suggested that theta activity may be associated with positive emotional responses to music.

The authors hypothesize that listening to K448 for as little as 30 seconds may activate networks within the brain that are associated with positive emotional responses to music and are regulated by the frontal cortex. The activation of these networks may contribute to reductions in epilepsy-associated [electrical activity](#) spikes among those with medication-resistant epilepsy, they suggest.

More information: Musical components important for the Mozart K448 effect in epilepsy, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-95922-7](https://doi.org/10.1038/s41598-021-95922-7), www.nature.com/articles/s41598-021-95922-7

Provided by Nature Publishing Group

APA citation: Therapeutic potential of Mozart for medication-resistant epilepsy (2021, September 16) retrieved 29 November 2021 from <https://medicalxpress.com/news/2021-09-therapeutic-potential-mozart-medication-resistant-epilepsy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.