

New research 'sniffs out' how associative memories are formed

September 22 2021



Credit: CC0 Public Domain

Has the scent of freshly baked chocolate chip cookies ever taken you back to afternoons at your grandmother's house? Has an old song ever brought back memories of a first date? The ability to remember

relationships between unrelated items (an odor and a location, a song and an event) is known as associative memory.

Psychologists began studying [associative memory](#) in the 1800s, with William James describing the phenomenon in his 1890 classic *The Principles of Psychology*. Scientists today agree that the structures responsible for the formation of associative [memory](#) are found in the medial temporal lobe, or the famous "memory center" of the brain, but the particular cells involved, and how those cells are controlled, have remained a mystery until now.

Neuroscientists at the University of California, Irvine have discovered specific types of neurons within the memory center of the brain that are responsible for acquiring new associative memories. Additionally, they have discovered how these associative memory neurons are controlled. We rely on associative memories in our [everyday lives](#) and this research is an important step in understanding the detailed mechanism of how these types of memories are formed in the brain.

"Although associative memory is one of the most basic forms of memory in our [everyday life](#), mechanisms underlying associative memory remain unclear" said lead researcher Kei Igarashi, faculty fellow of the Center for the Neurobiology of Learning and Memory and assistant professor of anatomy & neurobiology at the UCI School of Medicine.

The study published today in the journal *Nature*, reports for the first time, that specific cells in the lateral entorhinal cortex of the [medial temporal lobe](#), called fan cells, are required for the acquisition of new associative memories and that these cells are controlled by dopamine, a brain chemical known to be involved in our experience of pleasure or reward.

In the study, researchers used electrophysiological recordings and optogenetics to record and control activity from fan cells in mice as they learn to associate specific odors with rewards. This approach led researchers to discover that fan cells compute and represent the association of the two new unrelated items (odor and reward). These fan cells are required for successful acquisition of new associative memories. Without these [cells](#), pre-learned associations can be retrieved, but the new associations cannot be acquired. Additionally acquiring new associations also requires dopamine.

"We never expected that dopamine is involved in the memory circuit. However, when the evidence accumulated, it gradually became clear that dopamine is involved," said Igarashi. "These experiments were like a detective story for us, and we are excited about the results."

This discovery is an important piece in the puzzle of understanding how memories are formed in the brain and lays a foundation on which other researchers can continue to build. Associative memory abilities are known to decline in neurodegenerative diseases like Alzheimer's Disease. Understanding the neurobiological mechanism of how these memories are formed is the first step to developing therapeutics to slow the loss of associative memory abilities in Alzheimer's Disease.

More information: Lee, J.Y. et al. Dopamine facilitates associative memory encoding in the entorhinal cortex. *Nature* (2021).
doi.org/10.1038/s41586-021-03948-8

Provided by University of California, Irvine

Citation: New research 'sniffs out' how associative memories are formed (2021, September 22) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2021-09-associative-memories.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.