

High selenium levels found in bird eggs

25 December 2006

Analysis of the eggs of wild birds in California's San Joaquin Valley appears to show that recycling agricultural runoff is causing high levels of selenium.

The U.S. Bureau of Reclamation is considering expansion of the recycling program, the Los Angeles Times reported. But this year's sampling of eggs found the highest levels of selenium in five years.

The drainage project keeps selenium-tainted water out of the San Joaquin River by using it to irrigate crops that can tolerate high levels of salt. Irrigation builds up levels of selenium and other chemicals that occur naturally by washing them out of the soil.

In the 1980s, water in farm drainage ponds became so toxic that it caused birth defects in birds.

"They have to be extremely careful; that water is so potent, they can't make any mistakes," said U.S. Fish and Wildlife biologist Joseph Skorupa. "If they make a mistake, they'll have dead birds."

Joe McGahan, coordinator of the reclamation program, said the high selenium levels could be a one-year spike. He said scientists have found no evidence that birds are dying or being born deformed.

Copyright 2006 by United Press International

APA citation: High selenium levels found in bird eggs (2006, December 25) retrieved 11 May 2021 from <https://medicalxpress.com/news/2006-12-high-selenium-bird-eggs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.